

## Product datasheet for **TA321763**


### **C13orf24 (PIBF1) Rabbit Polyclonal Antibody**

#### **Product data:**


Product Type:	Primary Antibodies
Applications:	IHC
Recommended Dilution:	IHC: 50-200 Positive control: Human colon cancer Predicted cell location: Cytoplasm
Reactivity:	Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Synthetic peptide peptide corresponding to a region derived from 745-757 amino acids of human progesterone immunomodulatory binding factor 1
Formulation:	PBS pH7.3, 0.05% NaN <sub>3</sub> , 50% glycerol
Concentration:	lot specific
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	progesterone immunomodulatory binding factor 1
Database Link:	<a href="#">NP_006337</a> <a href="#">Entrez Gene 10464 Human</a> <a href="#">Q8WXW3</a>
Background:	Progesterone-induced-blocking factor 1 is a protein that in humans is encoded by the PIBF1 gene. It has been shown to localize to the centrosome and has also been named CEP90. Mediator of progesterone that by acting on the phospholipase A2 enzyme interferes with arachidonic acid metabolism; induces a Th2 biased immune response; and by controlling NK activity exerts an anti-abortive effect.
Synonyms:	C13orf24; CEP90; PIBF




[View online »](#)

**Product images:**


Immunohistochemistry of paraffin-embedded Human colon cancer tissue using TA321763 (PIBF1 Antibody) at dilution 1/60 (Original magnification:  $\times 200$ )



Immunohistochemistry of paraffin-embedded Human colon cancer tissue using TA321763 (PIBF1 Antibody) at dilution 1/60, treated with synthetic peptide. (Original magnification:  $\times 200$ )



Immunohistochemistry of paraffin-embedded Human lung cancer tissue using TA321763 (PIBF1 Antibody) at dilution 1/60 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human lung cancer tissue using TA321763 (PIBF1 Antibody) at dilution 1/60, treated with synthetic peptide. (Original magnification: ×200)