

Product datasheet for TA321793

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Glucose Transporter GLUT3 (SLC2A3) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 500-2000

WB positive control: Human colon cancer tissue

Reactivity: Human
Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide corresponding to a region derived from 466-479 amino acids of human

solute carrier family 2 (facilitated glucose transporter), member 3

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 54 kDa

Gene Name: solute carrier family 2 member 3

Database Link: NP 008862

Entrez Gene 6515 Human

P11169



Background:

Glucose transporter 3?(or?GLUT3); also known as?solute carrier family 2; facilitated glucose transporter member 3?(SLC2A3) is a?protein?that in humans is encoded by the?SLC2A3?gene. GLUT3?facilitates?the transport ofglucose?across the?plasma membranes?of mammalian cells. GLUT3 is most known for its specific expression in?neurons?and has originally been designated as the neuronal GLUT. GLUT3 has been studied in other cell types with specific glucose requirements; including?sperm; preimplantation?embryos; circulating?white blood cells?and?carcinoma?cell lines. GLUT3 has both a higher affinity for glucose and at least a fivefold greater transport capacity than GLUT1; GLUT2 and GLUT4; which is particularly significant for its role in neuronal glucose transport; where ambient glucose levels are fivefold lower than in serum.

Synonyms: GLUT3

Protein Families: Transmembrane

Product images:



Gel: 10%SDS-PAGE Lysate: 40 μg

Lane: Human colon cancer tissue

Primary antibody: TA321793 (SLC2A3 Antibody) at

dilution 1/400

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 2 minutes