

Product datasheet for TA321811

CLCA4 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 1000-5000

WB positive control: Rat esophagus tissue lysate

IHC: 50-200

Positive control: Human thyroid cancer

Predicted cell location: Cytoplasm and Cell membrane

Reactivity: Human, Rat

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein corresponding to a region derived from 306-476 amino acids of Human

Calcium-activated chloride channel regulator 4

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 101 kDa

Gene Name: chloride channel accessory 4

Database Link: NP 036260

Entrez Gene 362053 RatEntrez Gene 22802 Human

Q14CN2



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background: The protein encoded by this gene belongs to the calcium sensitive chloride conductance

protein family. To date; all members of this gene family map to the same site on

chromosome 1p31-p22 and share high degrees of homology in size; sequence and predicted structure; but differ significantly in their tissue distributions. Alternative splicing results in


multiple transcript variants; only one of which is thought to be protein coding.


Synonyms: CaCC; CaCC2

Protein Families: Druggable Genome, Ion Channels: Other, Transmembrane

Protein Pathways: Olfactory transduction

Product images:





Gel: 6%SDS-PAGE Lysate: 40 μg

Lane: Rat esophagus tissue lysate

Primary antibody: TA321811 (CLCA4 Antibody) at

dilution 1/500

Secondary antibody: Goat anti rabbit IgG at

1/5000 dilution

Exposure time: 30 seconds

Primary antibody: TA321811 (CLCA4 Antibody) at

dilution 1/500


Secondary antibody: Goat anti rabbit IgG at

1/5000 dilution


Exposure time: 30 seconds

Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA321811 (CLCA4 Antibody) at dilution 1/70 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA321811 (CLCA4 Antibody) at dilution 1/70, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA321811 (CLCA4 Antibody) at dilution 1/70 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA321811 (CLCA4 Antibody) at dilution 1/70, treated with fusion protein. (Original magnification: ×200)