

Product datasheet for TA321939

CHRFAM7A Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Mouse heart tissue

IHC: 25-100

Positive control: Human liver cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide peptide corresponding to a region derived from 32-46 amino acids of

human CHRNA7 (cholinergic receptor, nicotinic, alpha 7, exons 5-10) and FAM7A (family with

sequence similarity 7A, exons A-E) fusion

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 46 kDa

Gene Name: CHRNA7 (exons 5-10) and FAM7A (exons A-E) fusion

Database Link: NP 683709

Entrez Gene 89832 Human

Q494W8



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:

The nicotinic acetylcholine receptors (nAChRs) are members of a superfamily of ligand-gated ion channels that mediate fast signal transmission at synapses. The family member CHRNA7; which is located on chromosome 15 in a region associated with several neuropsychiatric disorders; is partially duplicated and forms a hybrid with a novel gene from the family with sequence similarity 7 (FAM7A). Alternative splicing has been observed; and two variants exist; for this hybrid gene. The N-terminally truncated products predicted by the largest open reading frames for each variant would lack the majority of the neurotransmitter-gated ion-channel ligand binding domain but retain the transmembrane region that forms the ion channel. Although current evidence supports transcription of this hybrid gene; translation of the nicotinic acetylcholine receptor-like protein-encoding open reading frames has not been confirmed.

Synonyms: CHRNA7; CHRNA7-DR1; D-10

Protein Families: Druggable Genome, Ion Channels: Other, Transmembrane

Product images:



Gel: 10%SDS-PAGE Lysate: 40 μg

Lane: Mouse heart tissue

Primary antibody: TA321939 (CHRFAM7A


Antibody) at dilution 1/700

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 2 minutes






Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA321939 (CHRFAM7A Antibody) at dilution 1/40 (Original magnification: ×200)




Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA321939 (CHRFAM7A Antibody) at dilution 1/40, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA321939 (CHRFAM7A Antibody) at dilution 1/40 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA321939 (CHRFAM7A Antibody) at dilution 1/40, treated with synthetic peptide. (Original magnification: ×200)