

## **Product datasheet for TA321945S**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

## **AGR3 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 1000-5000

WB positive control: A549 cell, HT-29 cell, Human next to colon cancer tissue lysates

IHC: 50-200

Positive control: Human thyroid cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

**Clonality:** Polyclonal

**Immunogen:** Fusion protein corresponding to a region derived from 22-166 amino acids of human anterior

gradient 3 homolog (Xenopus laevis)

**Formulation:** PBS pH7.3, 0.05% NaN3, 50% glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 19 kDa

**Gene Name:** anterior gradient 3, protein disulphide isomerase family member

Database Link: NP 789783

Entrez Gene 403205 MouseEntrez Gene 155465 Human

Q8TD06






Background:

Anterior gradient protein 3 homolog?is a?protein?that in humans is encoded by the?AGR3? gene. The deduced 166-amino acid protein has an N-terminal signal sequence. Fluorescence-tagged BCMP11 localized in secretory or endosome-like organelles in transfected breast carcinoma cell lines. Immunohistochemical analysis showed that expression of BCMP11 was restricted in normal tissues, with significant staining of only the epithelial lining of colonic mucosa. However, BCMP11 was detected in 43 of 58 (74%) breast cancer tissues tested, but not in normal breast ductal epithelial cells. In cancerous epithelial cells, BCMP11 localized to the cytoplasm.

Synonyms: AG-3; AG3; BCMP11; hAG-3; HAG3; PDIA18

**Protein Families:** Secreted Protein

## **Product images:**



Gel: 15%SDS-PAGE Lysate: 40 μg Lane 1-2: MCF7 cells human colon tissue


Primary antibody: [TA321945] (AGR3 Antibody) at

dilution 1/150

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 20 seconds




Immunohistochemistry of paraffin-embedded Human colon cancer tissue using [TA321945] (AGR3 Antibody) at dilution 1/15 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human colon cancer tissue using [TA321945] (AGR3 Antibody) at dilution 1/15, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA321945] (AGR3 Antibody) at dilution 1/15 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA321945] (AGR3 Antibody) at dilution 1/15, treated with fusion protein. (Original magnification: ×200)