

## **Product datasheet for TA321951S**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Telomerase reverse transcriptase (TERT) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Human mucinous type of soft tissue sarcoma tissue lysate

IHC: 100-300

Positive control: Human colorectal cancer Predicted cell location: Nucleus and Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

**Clonality:** Polyclonal

**Immunogen:** Synthetic peptide corresponding to a region derived from 1120-1132 amino acids of human

telomerase reverse transcriptase

**Formulation:** PBS pH7.3, 0.05% NaN3, 50% glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** telomerase reverse transcriptase

Database Link: NP 937986

Entrez Gene 7015 Human

014746





Background:

Telomerase is a ribonucleoprotein polymerase that maintains telomere ends by addition of the telomere repeat TTAGGG. The enzyme consists of a protein component with reverse transcriptase activity; encoded by this gene; and an RNA component which serves as a template for the telomere repeat. Telomerase expression plays a role in cellular senescence; as it is normally repressed in postnatal somatic cells resulting in progressive shortening of telomeres. Deregulation of telomerase expression in somatic cells may be involved in oncogenesis. Studies in mouse suggest that telomerase also participates in chromosomal repair; since de novo synthesis of telomere repeats may occur at double-stranded breaks. Alternatively spliced variants encoding different isoforms of telomerase reverse transcriptase have been identified; the full-length sequence of some variants has not been determined. Alternative splicing at this locus is thought to be one mechanism of regulation of telomerase activity.


Synonyms:

EST2; hEST2; TCS1; telomerase catalytic subunit; telomerase reverse transcriptase; TP2; TRT

**Protein Families:** 

Druggable Genome

## **Product images:**



Gel: 6%SDS-PAGE Lysate: 40 μg

Lane: Human mucinous type of soft tissue

sarcoma tissue lysate

Primary antibody: [TA321951] (TERT Antibody) at


dilution 1/700

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 90 seconds






Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using [TA321951] (TERT Antibody) at dilution 1/70 (Original magnification: ×200)




Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using [TA321951] (TERT Antibody) at dilution 1/70, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA321951] (TERT Antibody) at dilution 1/70 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA321951] (TERT Antibody) at dilution 1/70, treated with synthetic peptide. (Original magnification: ×200)