

Product datasheet for TA322014

SLC22A6 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Human fetal kidney tissue

IHC: 50-200

Positive control: Human colon cancer Predicted cell location: Cytoplasm

Reactivity: Human, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide corresponding to a region derived from 551-563 amino acids of Human

solute carrier family 22 (organic anion transporter), member 6

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 56 kDa

Gene Name: solute carrier family 22 member 6

Database Link: NP 695009

Entrez Gene 29509 RatEntrez Gene 9356 Human

Q4U2R8



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

The protein encoded by this gene is involved in the sodium-dependent transport and excretion of organic anions; some of which are potentially toxic. The encoded protein is an integral membrane protein and may be localized to the basolateral membrane. Four transcript variants encoding four different isoforms have been found for this gene. Functions as organic anion exchanger when the uptake of one molecule of organic anion is coupled with an efflux of one molecule of endogenous dicarboxylic acid (glutarate; ketoglutarate; etc). Mediates the sodium-independent uptake of 2;3-dimercapto-1-propanesulfonic acid (DMPS) By similarity.

Synonyms: HOAT1; OAT1; PAHT; ROAT1

Protein Families: Transmembrane

Product images:



Gel: 10%SDS-PAGE Lysate: 40 μg

Lane: Human fetal kidney tissue


Primary antibody: TA322014 (SLC22A6 Antibody)

at dilution 1/600

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 3 minutes




Immunohistochemistry of paraffin-embedded Human colon cancer tissue using TA322014 (SLC22A6 Antibody) at dilution 1/40 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human colon cancer tissue using TA322014 (SLC22A6 Antibody) at dilution 1/40, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA322014 (SLC22A6 Antibody) at dilution 1/40 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA322014 (SLC22A6 Antibody) at dilution 1/40, treated with synthetic peptide. (Original magnification: ×200)