

## **Product datasheet for TA322053S**

### 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc.

#### EU: info-de@origene.com CN: techsupport@origene.cn

## Kv3.2 (KCNC2) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB: 200-1000

WB positive control: 231 cells and human brain malignant glioma tissue

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Synthetic peptide corresponding to a region derived from 16-30 amino acids of Human

potassium voltage-gated channel, Shaw-related subfamily, member 2

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

**Purification:** Antigen affinity purification

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 62 kDa

**Gene Name:** potassium voltage-gated channel subfamily C member 2

Database Link: NP 631875

Entrez Gene 246153 RatEntrez Gene 268345 MouseEntrez Gene 3747 Human

Q96PR1

**Background:** The Shaker gene family of Drosophila encodes components of voltage-gated potassium

channels and is comprised of four subfamilies. Based on sequence similarity; this gene is similar to one of these subfamilies; namely the Shaw subfamily. The protein encoded by this gene belongs to the delayed rectifier class of channel proteins and is an integral membrane protein that mediates the voltage-dependent potassium ion permeability of excitable

membranes. Several transcript variants encoding different isoforms have been found for this

gene.

Synonyms: KV3.2





**Protein Families:** Druggable Genome, Ion Channels: Potassium, Transmembrane

# **Product images:**



Gel: 8%SDS-PAGE Lysate: 50 µg Lane 1-2: 231 cells

human brain malignant glioma tissue

Primary antibody: [TA322053] (KCNC2 Antibody)

at dilution 1/450

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 10 seconds