

## **Product datasheet for TA322061**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc.

EU: info-de@origene.com
CN: techsupport@origene.cn

## **ADAMTS17 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: ELISA: 1:2000-5000, WB: 1:500-2000, IHC: 1:25-100

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Synthetic peptide corresponding to a region derived from 609-621 amino acids of human

ADAM metallopeptidase with thrombospondin type 1 motif, 17

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 121 kDa

Gene Name: ADAM metallopeptidase with thrombospondin type 1 motif 17

Database Link: NP 620688

Entrez Gene 233332 MouseEntrez Gene 293004 RatEntrez Gene 170691 Human

**Q8TE56** 

**Background:** This gene encodes a member of the ADAMTS (a disintegrin and metalloproteinase with

thrombospondin motifs) protein family. ADAMTS family members share several distinct protein modules; including a propeptide region; a metalloproteinase domain; a disintegrin-like domain; and a thrombospondin type 1 (TS) motif. Individual members of this family differ in the number of C-terminal TS motifs; and some have unique C-terminal domains. The protein encoded by this gene has a high sequence similarity to the protein encoded by ADAMTS19; another family member. The function of this protein has not been determined.


**Synonyms:** FLJ16363; FLJ32769






**Protein Families:** Druggable Genome

## **Product images:**



Predicted band size: 121 kDa. Positive control: NIH/3T3 cell lysate. Recommended dilution: 1/500-2000. (Gel: 8%SDS-PAGE Lysate: 40 ug per lane Primary antibody: 1/1200 dilution Secondary antibody: Goat anti Rabbit lgG - H&L (HRP) at 1/10000 dilution Exposure time: 1 hour)



Predicted cell location: Cytoplasm. Positive control: Human colon cancer tissue.

Recommended dilution: 1/25-100 The image on the left is immunohistochemistry of paraffinembedded human colon cancer tissue using ADAMTS17 antibody at dilution 1/60, on the right is treated with the synthetic peptide. (Original magnification:x200)