

Product datasheet for TA322068S

SPATA17 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: Human liver cancer tissue

IHC: 25-100

Positive control: Human liver cancer Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein corresponding to C terminal 300 amino acids of human spermatogenesis

associated 17

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 43 kDa

Gene Name: spermatogenesis associated 17

Database Link: NP 620151

Entrez Gene 128153 Human

Q96L03



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Spata17?gene, which called MSRG11, by digital differential display of ESTs found only in testis. The deduced 292-amino acid protein was considered to be a member of the family of calmodulin (CaM)-binding proteins because it contained 3 short CaM-binding motifs containing conserved ile and gln residues (IQ motif). Spata17 protein was most abundant in the cytoplasm of round spermatids and elongating spermatids within seminiferous tubules of the adult testis. Expression gradually decreased in the manipulated testis of animals that underwent experimental unilateral cryptorchidism. Transient transfection experiments with a Spata17 expression construct in cultured spermatogonia GC-1 cells indicated that Spata17 accelerated apoptosis in a dose-dependent manner.

Synonyms:

IQCH; MSRG-11; MSRG11

Product images:



Gel: 10%SDS-PAGE Lysate: 40 μg


Lane: Human liver cancer tissue Primary antibody: [TA322068] (SPATA17

Antibody) at dilution 1/400

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 1 minute




Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA322068] (SPATA17 Antibody) at dilution 1/25 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA322068] (SPATA17 Antibody) at dilution 1/25, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human lung cancer tissue using [TA322068] (SPATA17 Antibody) at dilution 1/25 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human lung cancer tissue using [TA322068] (SPATA17 Antibody) at dilution 1/25, treated with fusion protein. (Original magnification: ×200)