

## **Product datasheet for TA322110**

## **Profilin 2 (PFN2) Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Human fetal brain tissue

IHC: 25-100

Positive control: Human brain Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

**Clonality:** Polyclonal

**Immunogen:** Synthetic peptide corresponding to a region derived from 128-140 amino acids of Human

profilin 2

**Formulation:** PBS pH7.3, 0.05% NaN3, 50% glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 15 kDa

Gene Name: profilin 2

Database Link: NP 444252

Entrez Gene 18645 MouseEntrez Gene 81531 RatEntrez Gene 5217 Human

P35080



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:

The protein encoded by this gene is a ubiquitous actin monomer-binding protein belonging to the profilin family. It is thought to regulate actin polymerization in response to extracellular signals. There are two alternatively spliced transcript variants encoding different isoforms described for this gene. Binds to actin and affects the structure of the cytoskeleton. At high concentrations; profilin prevents the polymerization of actin; whereas it enhances it at low concentrations. By binding to PIP2; it inhibits the formation of IP3 and DG.

Synonyms: D3S1319E; PFL

**Protein Families:** Druggable Genome

**Protein Pathways:** Regulation of actin cytoskeleton

## **Product images:**

KDa 40— 35— 25—

Gel: 10%SDS-PAGE Lysate: 40 μg

Lane: Human fetal brain tissue


Primary antibody: TA322110 (PFN2 Antibody) at

dilution 1/450

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 3 minutes




Immunohistochemistry of paraffin-embedded Human brain tissue using TA322110 (PFN2 Antibody) at dilution 1/30 (Original magnification: x200)






Immunohistochemistry of paraffin-embedded Human brain tissue using TA322110 (PFN2 Antibody) at dilution 1/30, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA322110 (PFN2 Antibody) at dilution 1/30 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA322110 (PFN2 Antibody) at dilution 1/30, treated with synthetic peptide. (Original magnification: ×200)