

Product datasheet for TA322155

Frizzled 8 (FZD8) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 50-200

Positive control: Human esophagus cancer

Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide corresponding to a region derived from 680-694 amino acids of human

frizzled family receptor 8

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: frizzled class receptor 8

Database Link: NP 114072

Entrez Gene 14370 MouseEntrez Gene 364754 RatEntrez Gene 8325 Human

Q9H461

Background: This intronless gene is a member of the frizzled gene family. Members of this family encode

seven-transmembrane domain proteins that are receptors for the Wingless type MMTV integration site family of signaling proteins. Most frizzled receptors are coupled to the beta-catenin canonical signaling pathway. This gene is highly expressed in two human cancer cell lines; indicating that it may play a role in several types of cancer. The crystal structure of the

extracellular cysteine-rich domain of a similar mouse protein has been determined.

Synonyms: FZ-8; hFZ8



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Frizzled 8 (FZD8) Rabbit Polyclonal Antibody - TA322155

Protein Families: Druggable Genome, Transmembrane


Protein Pathways: Basal cell carcinoma, Colorectal cancer, Melanogenesis, Pathways in cancer, Wnt signaling

pathway

Product images:




Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA322155 (FZD8 Antibody) at dilution 1/50 (Original magnification: ×200)




Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA322155 (FZD8 Antibody) at dilution 1/50, treated with synthetic peptide. (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA322155 (FZD8 Antibody) at dilution 1/50 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA322155 (FZD8 Antibody) at dilution 1/50, treated with synthetic peptide. (Original magnification: ×200)