

Product datasheet for TA322184S

NPPC Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 50-200

Positive control: Human gasrtic cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide corresponding to a region derived from 111-125 amino acids of Human

notch 4

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: natriuretic peptide C

Database Link: NP 077720

Entrez Gene 18159 MouseEntrez Gene 114593 RatEntrez Gene 4880 Human

P23582



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

This gene encodes a member of the Notch family. Members of this Type 1 transmembrane protein family share structural characteristics including an extracellular domain consisting of multiple epidermal growth factor-like (EGF) repeats; and an intracellular domain consisting of multiple; different domain types. Notch family members play a role in a variety of developmental processes by controlling cell fate decisions. The Notch signaling network is an evolutionarily conserved intercellular signaling pathway which regulates interactions between physically adjacent cells. In Drosophilia; notch interaction with its cell-bound ligands (delta; serrate) establishes an intercellular signaling pathway that plays a key role in development. Homologues of the notch-ligands have also been identified in human; but precise interactions between these ligands and the human notch homologues remain to be determined. This protein is cleaved in the trans-Golgi network; and presented on the cell surface as a heterodimer. This protein functions as a receptor for membrane bound ligands; and may play a role in vascular; renal and hepatic development. This gene may be associated with susceptibility to schizophrenia in a small portion of cases. An alternative splice variant has been described but its biological nature has not been determined.

Synonyms: CNP; CNP2


Protein Families: Secreted Protein

Product images:



Immunohistochemistry of paraffin-embedded Human gasrtic cancer tissue using [TA322184] (NPPC Antibody) at dilution 1/70 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human gasrtic cancer tissue using [TA322184] (NPPC Antibody) at dilution 1/70, treated with synthetic peptide. (Original magnification: ×200)