

## **Product datasheet for TA322202S**

## **Livin (BIRC7) Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Mouse adrenal gland tissue lysate

IHC: 50-200

Positive control: Human gastric cancer

Predicted cell location: Cytoplasm and Nucleus

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Full length fusion protein

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 33 kDa

**Gene Name:** baculoviral IAP repeat containing 7

Database Link: NP 071444

Entrez Gene 329581 MouseEntrez Gene 79444 Human

Q96CA5



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:


The protein encoded by this gene is a member of the family of inhibitor of apoptosis proteins (IAP) and contains a single copy of a baculovirus IAP repeat (BIR) as well as a RING-type zinc finger domain. The BIR domain is essential for inhibitory activity and interacts with caspases; while the RING finger domain sometimes enhances antiapoptotic activity but does not inhibit apoptosis alone. Two transcript variants encoding different isoforms have been found for this gene. The two isoforms have different antiapoptotic properties; with isoform alpha protecting cells from apoptosis induced by staurosporine and isoform b protecting cells from apoptosis induced by etoposide.?

Synonyms: KIAP; LIVIN; ML-IAP; MLIAP; RNF50

**Protein Families:** Druggable Genome

## **Product images:**








Lane: Mouse adrenal gland tissue lysate Primary antibody: [TA322202] (BIRC7 Antibody) at

dilution 1/700

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 90 seconds



Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using [TA322202] (BIRC7 Antibody) at dilution 1/75 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using [TA322202] (BIRC7 Antibody) at dilution 1/75, treated with fusion protein. (Original magnification: ×200)