

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA322255

GJC2 Rabbit Polyclonal Antibody

Product data:


| Due du et True et | |
|-------------------------|---|
| Product Type: | Primary Antibodies |
| Applications: | WB |
| Recommended Dilution: | WB: 500-2000 |
| | WB positive control: OP9 cells |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| lsotype: | IgG |
| Clonality: | Polyclonal |
| Immunogen: | Synthetic peptide corresponding to a region derived from 420-434 amino acids of Human gap junction protein, gamma 2, 47kDa |
| Formulation: | PBS pH7.3, 0.05% NaN3, 50% glycerol |
| Concentration: | lot specific |
| Purification: | Antigen affinity purification |
| Conjugation: | Unconjugated |
| Storage: | Store at -20°C as received. |
| Stability: | Stable for 12 months from date of receipt. |
| Predicted Protein Size: | 47 kDa |
| Gene Name: | gap junction protein gamma 2 |
| Database Link: | <u>NP_065168</u> <u>Entrez Gene 118454 MouseEntrez Gene 497913 RatEntrez Gene 57165 Human</u> <u>Q5T442</u> |
| Background: | This gene encodes a gap junction protein. Gap junction proteins are members of a large family of homologous connexins and comprise 4 transmembrane; 2 extracellular; and 3 cytoplasmic domains. This gene plays a key role in central myelination and is involved in peripheral myelination in humans. Defects in this gene are the cause of autosomal recessive Pelizaeus-Merzbacher-like disease-1. |
| Synonyms: | CX46.6; Cx47; GJA12; HLD2; LMPH1C; PMLDAR; SPG44 |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US Protein Families: Ic

Ion Channels: Other

Product images:



Gel: 10%SDS-PAGE Lysate: 30 µg Lane: OP9 cells Primary antibody: TA322255 (GJC2 Antibody) at dilution 1/450 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution Exposure time: 30 seconds

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US