

Product datasheet for TA322582S

NGAL (LCN2) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 25-100

Positive control: Human brain Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide peptide corresponding to a region derived from 60-73 amino acids of

human lipocalin 2

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: lipocalin 2

Database Link: NP 005555

Entrez Gene 3934 Human

P80188



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Iron-trafficking protein involved in multiple processes such as apoptosis; innate immunity and renal development. Binds iron through association with 2;5-dihydroxybenzoic acid (2;5-DHBA); a siderophore that shares structural similarities with bacterial enterobactin; and delivers or removes iron from the cell; depending on the context. Iron-bound form (holo-24p3) is internalized following binding to the SLC22A17 (24p3R) receptor; leading to release of iron and subsequent increase of intracellular iron concentration. In contrast; association of the iron-free form (apo-24p3) with the SLC22A17 (24p3R) receptor is followed by association with an intracellular siderophore; iron chelation and iron transfer to the extracellular medium; thereby reducing intracellular iron concentration. Involved in apoptosis due to interleukin-3 (IL3) deprivation: iron-loaded form increases intracellular iron concentration without promoting apoptosis; while iron-free form decreases intracellular iron levels; inducing expression of the proapoptotic protein BCL2L11/BIM; resulting in apoptosis. Involved in innate immunity; possibly by sequestrating iron; leading to limit bacterial growth.


Synonyms: 24p3; MSFI; NGAL Protein Families: Secreted Protein

Product images:




Immunohistochemistry of paraffin-embedded Human brain tissue using [TA322582] (LCN2 Antibody) at dilution 1/50 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human brain tissue using [TA322582] (LCN2 Antibody) at dilution 1/50, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA322582] (LCN2 Antibody) at dilution 1/50 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA322582] (LCN2 Antibody) at dilution 1/50, treated with synthetic peptide. (Original magnification: ×200)