

## **Product datasheet for TA322603S**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Acyl coenzyme A Thioesterase 8 (ACOT8) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: Human testis tissue lysate

IHC: 15-50

Positive control: Human cervical cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Full length fusion protein

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** acyl-CoA thioesterase 8

Database Link: NP 005460

Entrez Gene 170588 RatEntrez Gene 170789 MouseEntrez Gene 10005 Human

<u>014734</u>

**Background:** The protein encoded by this gene is a peroxisomal thioesterase that appears to be involved

more in the oxidation of fatty acids rather than in their formation. The encoded protein can bind to the human immunodeficiency virus-1 protein Nef, and mediate Nef-induced down-regulation of CD4 in T-cells.?Acyl-CoA thioesterases are a group of enzymes that catalyze the hydrolysis of acyl-CoAs to the free fatty acid and coenzyme A (CoASH), providing the potential to regulate intracellular levels of acyl-CoAs, free fatty acids and CoASH. May mediate Nef-

induced down-regulation of CD4. Major thioesterase in peroxisomes.






Synonyms:

hacte-III; hnaacte; hte; nap1; pte-1; pte-2; pte1; pte2

## **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 μg

Lane: Human testis tissue lysate


Primary antibody: [TA322603] (ACOT8 Antibody)

at dilution 1/200


Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution


Exposure time: 1 minute




Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA322603] (ACOT8 Antibody) at dilution 1/15 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA322603] (ACOT8 Antibody) at dilution 1/15, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human lung cancer tissue using [TA322603] (ACOT8 Antibody) at dilution 1/15 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human lung cancer tissue using [TA322603] (ACOT8 Antibody) at dilution 1/15, treated with fusion protein. (Original magnification: ×200)