

## **Product datasheet for TA322683S**

## **MRP1 (ABCC1) Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: LO2 cell

IHC: 25-100

Positive control: Human liver cancer

Predicted cell location: Cytoplasm and Cell membrane

Reactivity: Human
Host: Rabbit
Isotype: IgG

**Clonality:** Polyclonal

**Immunogen:** Synthetic peptide corresponding to a region derived from 625-638 amino acids of human

ATP-binding cassette, sub-family C (CFTR/MRP), member 1

**Formulation:** PBS pH7.3, 0.05% NaN3, 50% glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** ATP binding cassette subfamily C member 1

Database Link: NP 004987

Entrez Gene 4363 Human

P33527



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:


The protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra-and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1; MDR/TAP; MRP; ALD; OABP; GCN20; White). This full transporter is a member of the MRP subfamily which is involved in multi-drug resistance. This protein functions as a multispecific organic anion transporter; with oxidized glutatione; cysteinyl leukotrienes; and activated aflatoxin B1 as substrates. This protein also transports glucuronides and sulfate conjugates of steroid hormones and bile salts. Alternatively spliced variants of this gene have been described but their full-length nature is unknown.

**Synonyms:** ABC29; ABCC; GS-X; MRP; MRP1

**Protein Families:** Druggable Genome, Transmembrane

**Protein Pathways:** ABC transporters

## **Product images:**



Gel: 6%SDS-PAGE Lysate: 40 μg Lane: LO2 cell

Primary antibody: [TA322683] (ABCC1 Antibody)


at dilution 1/200

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 30 seconds





Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA322683] (ABCC1 Antibody) at dilution 1/25 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA322683] (ABCC1 Antibody) at dilution 1/25, treated with synthetic peptide. (Original magnification: ×200)