

Product datasheet for TA322700S

ROCK2 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Mouse brain tissue lysate

IHC: 50-200

Positive control: Human tonsil Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein corresponding to C terminal 300 amino acids of human Rho-associated,

coiled-coil containing protein kinase 2

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 161 kDa

Gene Name: Rho associated coiled-coil containing protein kinase 2

Database Link: NP 004841

Entrez Gene 19878 MouseEntrez Gene 25537 RatEntrez Gene 9475 Human

<u>075116</u>



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background: The protein encoded by this gene is a serine/threonine kinase that regulates cytokinesis;

smooth muscle contraction; the formation of actin stress fibers and focal adhesions; and the activation of the c-fos serum response element. This protein; which is an isozyme of ROCK1 is

a target for the small GTPase Rho. Protein kinase which is a key regulator of actin

cytoskeleton and cell polarity. Involved in regulation of smooth muscle contraction; actin cytoskeleton organization; stress fiber and focal adhesion formation; neurite retraction; cell adhesion and motility via phosphorylation of ADD1; BRCA2; CNN1; EZR; DPYSL2; EP300; MSN;

MYL9/MLC2; NPM1; RDX; PPP1R12A and VIM. Phosphorylates SORL1 and IRF4.

Synonyms: ROCK-II


Protein Families: Druggable Genome, Protein Kinase

Protein Pathways: Axon guidance, Chemokine signaling pathway, Focal adhesion, Leukocyte transendothelial

migration, Pathogenic Escherichia coli infection, Regulation of actin cytoskeleton, TGF-beta

signaling pathway, Vascular smooth muscle contraction, Wnt signaling pathway

Product images:



Gel: 8%SDS-PAGE Lysate: 40 µg

Lane: Mouse brain tissue lysate

Primary antibody: [TA322700] (ROCK2 Antibody)


at dilution 1/800

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 2 minutes





Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA322700] (ROCK2 Antibody) at dilution 1/130 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA322700] (ROCK2 Antibody) at dilution 1/130, treated with fusion protein. (Original magnification: ×200)