

Product datasheet for TA322731S

LRP2 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 50-200

Positive control: Human esophagus cancer

Predicted cell location: Cell membrane, Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide corresponding to a region derived from 4640-4054 amino acids of Human

low density lipoprotein receptor-related protein 2

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: LDL receptor related protein 2

Database Link: NP 004516

Entrez Gene 14725 MouseEntrez Gene 29216 RatEntrez Gene 4036 Human

P98164



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:


The protein encoded by this gene; low density lipoprotein-related protein 2 (LRP2) or megalin; is a multi-ligand endocytic receptor that is expressed in many different tissues but primarily in absorptive epithilial tissues such as the kidney. This glycoprotein has a large aminoterminal extracellular domain; a single transmembrane domain; and a short carboxy-terminal cytoplasmic tail. The extracellular ligand-binding-domains bind diverse macromolecules including albumin; apolipoproteins B and E; and lipoprotein lipase. The LRP2 protein is critical for the reuptake of numerous ligands; including lipoproteins; sterols; vitamin-binding proteins; and hormones. This protein also has a role in cell-signaling; extracellular ligands include parathyroid horomones and the morphogen sonic hedgehog while cytosolic ligands include MAP kinase scaffold proteins and JNK interacting proteins. Recycling of this membrane receptor is regulated by phosphorylation of its cytoplasmic domain. Mutations in this gene cause Donnai-Barrow syndrome (DBS) and facio-oculoacoustico-renal syndrome (FOAR).

Synonyms: DBS; GP330

Protein Families: Druggable Genome, Transmembrane


Protein Pathways: Hedgehog signaling pathway

Product images:



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using [TA322731] (LRP2 Antibody) at dilution 1/40 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using [TA322731] (LRP2 Antibody) at dilution 1/40, treated with synthetic peptide. (Original magnification: ×200)