

Product datasheet for TA322828

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Vesicle docking protein p115 (USO1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Human fetal kidney tissue

IHC: 25-100

Positive control: Human ovarian cancer Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide corresponding to a region derived from 950-961 amino acids of human

USO1 vesicle transport factor

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 108 kDa

Gene Name: USO1 vesicle transport factor

Database Link: NP 003706

Entrez Gene 8615 Human

<u>060763</u>





Background:

The protein encoded by this gene is a peripheral membrane protein which recycles between the cytosol and the Golgi apparatus during interphase. It is regulated by phosphorylation: dephosphorylated protein associates with the Golgi membrane and dissociates from the membrane upon phosphorylation. Ras-associated protein 1 recruits this protein to coat protein complex II (COPII) vesicles during budding from the endoplasmic reticulum; where it interacts with a set of COPII vesicle-associated SNAREs to form a cis-SNARE complex that promotes targeting to the Golgi apparatus. Transport from the ER to the cis/medial Golgi compartments requires the action of this gene product; GM130 and giantin in a sequential manner.

Synonyms: P115; TAP; VDP

Product images:



Gel: 8%SDS-PAGE Lysate: 40 μg

Lane: Human fetal kidney tissue


Primary antibody: TA322828 (USO1 Antibody) at

dilution 1/400

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 5 seconds



Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using TA322828 (USO1 Antibody) at dilution 1/13 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using TA322828 (USO1 Antibody) at dilution 1/13, treated with synthetic peptide. (Original magnification: ×200)