

Product datasheet for TA322853

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

14-3-3 beta (YWHAB) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WE

Recommended Dilution: WB: 500-2000

WB positive control: Hela cells

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide corresponding to a region derived from 72-87 amino acids of Human

tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein, beta

polypeptide

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 28 kDa

Gene Name: tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein beta

Database Link: NP 003395

Entrez Gene 54401 MouseEntrez Gene 56011 RatEntrez Gene 7529 Human

P31946

Background: This gene encodes a protein belonging to the 14-3-3 family of proteins; members of which

mediate signal transduction by binding to phosphoserine-containing proteins. This highly conserved protein family is found in both plants and mammals. The encoded protein has been shown to interact with RAF1 and CDC25 phosphatases; suggesting that it may play a role in linking mitogenic signaling and the cell cycle machinery. Two transcript variants; which

encode the same protein; have been identified for this gene.






Synonyms: GW128; HEL-S-1; HS1; KCIP-1; YWHAA

Protein Families: Druggable Genome

Protein Pathways: Cell cycle, Neurotrophin signaling pathway, Oocyte meiosis

Product images:



Gel: 12%SDS-PAGE Lysate: 30 µg Lane: Hela cells

Primary antibody: TA322853 (YWHAB Antibody)

at dilution 1/550

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 5 seconds