

Product datasheet for TA322864

TXN Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 1000-5000

WB positive control: Hela cells and human liver cancer tissue

IHC: 50-200

Positive control: Human liver cancer Predicted cell location: Cytoplasm

Reactivity: Human, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide corresponding to a region derived from 90-105 amino acids of human

thioredoxin

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 12 kDa

Gene Name: thioredoxin

Database Link: NP 003320

Entrez Gene 7295 Human

P10599



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

The protein encoded by this gene acts as a homodimer and is involved in many redox reactions. The encoded protein is active in the reversible S-nitrosylation of cysteines in certain proteins; which is part of the response to intracellular nitric oxide. This protein is found in the cytoplasm. Two transcript variants encoding different isoforms have been found for this gene. Plays a role in the reversible S-nitrosylation of cysteine residues in target proteins; and thereby contributes to the response to intracellular nitric oxide. Nitrosylates the active site Cys of CASP3 in response to nitric oxide (NO); and thereby inhibits caspase-3 activity.

Synonyms: TRDX; TRX1

Protein Families: Druggable Genome

Product images:



Gel: 12%SDS-PAGE Lysate: 40 µg Lane 1-2: Hela cells human liver cancer tissue


Primary antibody: TA322864 (TXN Antibody) at

dilution 1/700

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 20 seconds




Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA322864 (TXN Antibody) at dilution 1/40 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA322864 (TXN Antibody) at dilution 1/40, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human lung cancer tissue using TA322864 (TXN Antibody) at dilution 1/40 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human lung cancer tissue using TA322864 (TXN Antibody) at dilution 1/40, treated with synthetic peptide. (Original magnification: ×200)