

Product datasheet for TA322968

PRKACG Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 25-100

Positive control: Human thyroid cancer Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein corresponding to a region derived from 44-298 amino acids of human protein

kinase, cAMP-dependent, catalytic, gamma

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: protein kinase cAMP-activated catalytic subunit gamma

Database Link: NP 002723

Entrez Gene 5568 Human

P22612

Background: cAMP-dependent protein kinase catalytic subunit gamma is an enzyme that in humans is

encoded by the PRKACG gene. Cyclic AMP-dependent protein kinase (PKA) consists of two catalytic subunits and a regulatory subunit dimer. This gene encodes the gamma form of its catalytic subunit. The gene is intronless and is thought to be a retrotransposon derived from the gene for the alpha form of the PKA catalytic subunit. PRKACG has been shown to interact

with Ryanodine receptor 2.

Synonyms: BDPLT19; KAPG; PKACg



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Protein Families: Druggable Genome, Protein Kinase


Protein Pathways: Apoptosis, Calcium signaling pathway, Chemokine signaling pathway, Dilated

cardiomyopathy, Gap junction, GnRH signaling pathway, Hedgehog signaling pathway, Insuling pathway, Long term potentiation, MARK signaling pathway, Molangaposis


signaling pathway, Long-term potentiation, MAPK signaling pathway, Melanogenesis, Olfactory transduction, Oocyte meiosis, Prion diseases, Progesterone-mediated oocyte maturation, Taste transduction, Vascular smooth muscle contraction, Vibrio cholerae

infection, Wnt signaling pathway

Product images:




Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA322968 (PRKACG Antibody) at dilution 1/25 (Original magnification: ×200)




Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA322968 (PRKACG Antibody) at dilution 1/25, treated with fusion protein. (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human lung cancer tissue using TA322968 (PRKACG Antibody) at dilution 1/25 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human lung cancer tissue using TA322968 (PRKACG Antibody) at dilution 1/25, treated with fusion protein. (Original magnification: ×200)