

## **Product datasheet for TA323003**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## p75 NGF Receptor (NGFR) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: ELISA: 1:2000-10000, WB: 1:500-2000, IHC: 1:50-200

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Synthetic peptide corresponding to a region derived from 188-211 amino acids of Human

nerve growth factor receptor

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 45 kDa

**Gene Name:** nerve growth factor receptor

Database Link: NP 002498

Entrez Gene 18053 MouseEntrez Gene 24596 RatEntrez Gene 4804 Human

P08138

**Background:** Nerve growth factor receptor contains an extracellular domain containing four 40-amino acid

repeats with 6 cysteine residues at conserved positions followed by a serine/threonine-rich region; a single transmembrane domain; and a 155-amino acid cytoplasmic domain. The

cysteine-rich region contains the nerve growth factor binding domain.

Synonyms: CD271; Gp80-LNGFR; p75(NTR); p75NTR; TNFRSF16


**Protein Families:** Druggable Genome, Transmembrane

**Protein Pathways:** Cytokine-cytokine receptor interaction, Neurotrophin signaling pathway






## **Product images:**



Predicted band size: 45 kDa. Positive control: Mouse heart tissue lysate. Recommended dilution: 1/500-2000. (Gel: 10%SDS-PAGE Lysate: 50 ug per lane Primary antibody: 1/950 dilution Secondary antibody: Goat anti Rabbit IgG - H&L (HRP) at 1/10000 dilution Exposure time: 20 seconds)



Predicted cell location: Cytoplasm. Positive control: Human tonsil tissue. Recommended dilution: 1/50-200 The image on the left is immunohistochemistry of paraffin-embedded human tonsil tissue using NGFR antibody at dilution 1/50, on the right is treated with the synthetic peptide. (Original magnification:x200)