

## **Product datasheet for TA323157**

## **HDAC2** Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 15-50

Positive control: Human breast cancer

Predicted cell location: Nucleus

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Synthetic peptide corresponding to a region derived from 474-488 amino acids of Human

histone deacetylase 2

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** histone deacetylase 2

Database Link: NP 001518

Entrez Gene 15182 MouseEntrez Gene 3066 Human

Q92769

**Background:** This gene product belongs to the histone deacetylase family. Histone deacetylases act via the

formation of large multiprotein complexes; and are responsible for the deacetylation of lysine residues at the N-terminal regions of core histones (H2A; H2B; H3 and H4). This protein forms transcriptional repressor complexes by associating with many different proteins; including YY1; a mammalian zinc-finger transcription factor. Thus; it plays an important role in transcriptional regulation; cell cycle progression and developmental events. Alternative

splicing results in multiple transcript variants.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms: HD2; RPD3; YAF1

**Protein Families:** Druggable Genome, Stem cell - Pluripotency, Transcription Factors


**Protein Pathways:** Cell cycle, Chronic myeloid leukemia, Huntington's disease, Notch signaling pathway,

Pathways in cancer

## **Product images:**




Immunohistochemistry of paraffin-embedded Human breast cancer tissue using TA323157 (HDAC2 Antibody) at dilution 1/17 (Original magnification: ×200)




Immunohistochemistry of paraffin-embedded Human breast cancer tissue using TA323157 (HDAC2 Antibody) at dilution 1/17, treated with synthetic peptide. (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human colon cancer tissue using TA323157 (HDAC2 Antibody) at dilution 1/17 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human colon cancer tissue using TA323157 (HDAC2 Antibody) at dilution 1/17, treated with synthetic peptide. (Original magnification: ×200)