

Product datasheet for TA323158S

HDAC2 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 25-100

Positive control: Human ovarian cancer

Predicted cell location: Nucleus

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide corresponding to a region derived from 474-488 amino acids of Human

histone deacetylase 2

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: histone deacetylase 2

Database Link: NP 001518

Entrez Gene 15182 MouseEntrez Gene 3066 Human

Q92769

Background: This gene product belongs to the histone deacetylase family. Histone deacetylases act via the

formation of large multiprotein complexes; and are responsible for the deacetylation of lysine residues at the N-terminal regions of core histones (H2A; H2B; H3 and H4). This protein forms transcriptional repressor complexes by associating with many different proteins; including YY1; a mammalian zinc-finger transcription factor. Thus; it plays an important role

in transcriptional regulation; cell cycle progression and developmental events. Alternative

splicing results in multiple transcript variants.

Synonyms: HD2; RPD3; YAF1



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Protein Families: Druggable Genome, Stem cell - Pluripotency, Transcription Factors


Protein Pathways: Cell cycle, Chronic myeloid leukemia, Huntington's disease, Notch signaling pathway,

Pathways in cancer

Product images:




Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using [TA323158] (HDAC2 Antibody) at dilution 1/25 (Original magnification: ×200)




Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using [TA323158] (HDAC2 Antibody) at dilution 1/25, treated with synthetic peptide. (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human colon cancer tissue using [TA323158] (HDAC2 Antibody) at dilution 1/25 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human colon cancer tissue using [TA323158] (HDAC2 Antibody) at dilution 1/25, treated with synthetic peptide. (Original magnification: ×200)