

# Product datasheet for TA323174

# **CELSR2** Rabbit Polyclonal Antibody

# **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	IHC
Recommended Dilution:	IHC: 50-200 Positive control: Human colon cancer Predicted cell location: Cytoplasm, Cell membrane
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
lsotype:	lgG
Clonality:	Polyclonal
Immunogen:	Synthetic peptide corresponding to a region derived from 2623-2639 amino acids of human cadherin, EGF LAG seven-pass G-type receptor 2
Formulation:	PBS pH7.3, 0.05% NaN3, 50% glycerol
Concentration:	lot specific
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	cadherin EGF LAG seven-pass G-type receptor 2
Database Link:	<u>NP_001399</u> <u>Entrez Gene 53883 MouseEntrez Gene 83465 RatEntrez Gene 1952 Human</u> <u>Q9HCU4</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


## **CELSR2** Rabbit Polyclonal Antibody – TA323174

**Background:** The protein encoded by this gene is a member of the flamingo subfamily; part of the cadherin superfamily. The flamingo subfamily consists of nonclassic-type cadherins; a subpopulation that does not interact with catenins. The flamingo cadherins are located at the plasma membrane and have nine cadherin domains; seven epidermal growth factor-like repeats and two laminin A G-type repeats in their ectodomain. They also have seven transmembrane domains; a characteristic unique to this subfamily. It is postulated that these proteins are receptors involved in contact-mediated communication; with cadherin domains acting as homophilic binding regions and the EGF-like domains involved in cell adhesion and receptor-ligand interactions. The specific function of this particular member has not been determined.

Synonyms: ADGRC2; CDHF10; EGFL2; Flamingo1; MEGF3


Protein Families: Druggable Genome, Transmembrane

## **Product images:**



Immunohistochemistry of paraffin-embedded Human colon cancer tissue using TA323174 (CELSR2 Antibody) at dilution 1/50 (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human colon cancer tissue using TA323174 (CELSR2 Antibody) at dilution 1/50, treated with synthetic peptide. (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US 

Immunohistochemistry of paraffin-embedded Human lung cancer tissue using TA323174 (CELSR2 Antibody) at dilution 1/50 (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human lung cancer tissue using TA323174 (CELSR2 Antibody) at dilution 1/50, treated with synthetic peptide. (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US