

Product datasheet for TA323178

EDA Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 100-500

WB positive control: Human lung cancer tissue

Reactivity: Human, Mouse

Host: Rabbit **IgG** Isotype:

Clonality: Polyclonal

Synthetic peptide corresponding to a region derived from 284-298 amino acids of Human Immunogen:

Ectodysplasin-A

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Store at -20°C as received. Storage:

Stable for 12 months from date of receipt. Stability:

Predicted Protein Size: 49 kDa

Gene Name: ectodysplasin A Database Link: NP 001390

Entrez Gene 13607 MouseEntrez Gene 1896 Human

Q92838

Background: The protein encoded by this gene is a type II membrane protein that can be cleaved by furin

to produce a secreted form. The encoded protein; which belongs to the tumor necrosis factor

family; acts as a homotrimer and may be involved in cell-cell signaling during the

development of ectodermal organs. Defects in this gene are a cause of ectodermal dysplasia;

anhidrotic; which is also known as X-linked hypohidrotic ectodermal dysplasia. Several transcript variants encoding many different isoforms have been found for this gene.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms: ECTD1; ED1-A1; ED1-A2; EDA-A1; EDA-A2; EDA1; EDA2; HED; HED1; ODT1; STHAGX1;

XHED; XLHED

Protein Families: Druggable Genome, Secreted Protein, Transmembrane

Protein Pathways: Cytokine-cytokine receptor interaction

Product images:



Gel: 10%SDS-PAGE Lysate: 30 µg

Lane: Human lung cancer tissue

Primary antibody: TA323178 (EDA Antibody) at

dilution 1/130

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 5 minutes