

Product datasheet for TA323179

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

AKR1C1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Human liver cancer and breast infiltrative duct tissue, human fetal brain

tissue IHC: 50-200

Positive control: Human colon cancer Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide corresponding to a region derived from 290-304 amino acids of human

aldo-keto reductase family 1, member C1

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 37 kDa

Gene Name: aldo-keto reductase family 1, member C1

Database Link: NP 001344

Entrez Gene 1645 Human

Q04828





Background:


This gene encodes a member of the aldo/keto reductase superfamily; which consists of more than 40 known enzymes and proteins. These enzymes catalyze the conversion of aldehydes and ketones to their corresponding alcohols by utilizing NADH and/or NADPH as cofactors. The enzymes display overlapping but distinct substrate specificity. This enzyme catalyzes the reaction of progesterone to the inactive form 20-alpha-hydroxy-progesterone. This gene shares high sequence identity with three other gene members and is clustered with those three genes at chromosome 10p15-p14.?

Synonyms: 2-ALPHA-HSD; 20-ALPHA-HSD; C9; DD1; DD2; DDH; DDH1; H-37; HAKRC; HBAB; MBAB

Protein Families: Druggable Genome

Protein Pathways: Metabolism of xenobiotics by cytochrome P450

Product images:



Gel: 10%SDS-PAGE Lysate: 27 μg

Lane 1-3: Human liver cancer tissue Human breast infiltrative duct tissue

human fetal brain tissue


Primary antibody: TA323179 (AKR1C1 Antibody)

at dilution 1/500

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 2 minutes




Immunohistochemistry of paraffin-embedded Human colon cancer tissue using TA323179 (AKR1C1 Antibody) at dilution 1/30 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human colon cancer tissue using TA323179 (AKR1C1 Antibody) at dilution 1/30, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using TA323179 (AKR1C1 Antibody) at dilution 1/30 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using TA323179 (AKR1C1 Antibody) at dilution 1/30, treated with synthetic peptide. (Original magnification: ×200)