

## **Product datasheet for TA323285S**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Somatostatin Receptor 1 (SSTR1) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: 293T and hela cells

IHC: 50-200

Positive control: Human liver cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

**Clonality:** Polyclonal

**Immunogen:** Synthetic peptide corresponding to a region derived from 370-384 amino acids of human

somatostatin receptor 1

**Formulation:** PBS pH7.3, 0.05% NaN3, 50% glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 43 kDa

**Gene Name:** somatostatin receptor 1

Database Link: NP 001040

Entrez Gene 20605 MouseEntrez Gene 25033 RatEntrez Gene 6751 Human

P30872





**Background:** Somatostatins are peptide hormones that regulate diverse cellular functions such as

neurotransmission; cell proliferation; and endocrine signaling as well as inhibiting the release of many hormones and other secretory proteins. Somatostatin has two active forms of 14 and 28 amino acids. The biological effects of somatostatins are mediated by a family of G-protein coupled somatostatin receptors that are expressed in a tissue-specific manner. The protein encoded by this gene is a member of the superfamily of somatostatin receptors having seven transmembrane segments. Somatostatin receptors form homodimers and heterodimers with other members of the superfamily as well as with other G-protein coupled receptors and receptor tyrosine kinases. This somatostatin receptor has greater affinity for

somatostatin-14 than -28.


**Protein Families:** Druggable Genome, GPCR, Transmembrane

SRIF-2; SS-1-R; SS1-R; SS1R

**Protein Pathways:** Neuroactive ligand-receptor interaction

## **Product images:**

Synonyms:



Gel: 10%SDS-PAGE Lysate: 60 µg Lane 1-2: 293T cells

hela cells

Primary antibody: [TA323285] (SSTR1 Antibody) at


dilution 1/500

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 30 seconds






Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA323285] (SSTR1 Antibody) at dilution 1/40 (Original magnification: ×200)




Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA323285] (SSTR1 Antibody) at dilution 1/40, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using [TA323285] (SSTR1 Antibody) at dilution 1/40 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using [TA323285] (SSTR1 Antibody) at dilution 1/40, treated with synthetic peptide. (Original magnification: ×200)