

Product datasheet for TA323291

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

ABCE1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 500-2000

WB positive control: NIH/3T3 cells and human ovarian cancer tissue

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein corresponding to a region derived from 345-562 amino acids of human ATP-

binding cassette, sub-family E (OABP), member 1

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 67 kDa

Gene Name: ATP binding cassette subfamily E member 1

Database Link: NP 001035809

Entrez Gene 24015 MouseEntrez Gene 6059 Human

P61221






Background:

The protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1; MDR/TAP; MRP; ALD; OABP; GCN20; White). This protein is a member of the OABP subfamily. Alternatively referred to as the RNase L inhibitor; this protein functions to block the activity of ribonuclease L. Activation of ribonuclease L leads to inhibition of protein synthesis in the 2-5A/RNase L system; the central pathway for viral interferon action. Two transcript variants encoding the same protein have been found for this gene.

Synonyms: ABC38; OABP; RLI; RNASEL1; RNASELI; RNS4I

Protein Families: Druggable Genome

Product images:



Gel: 10%SDS-PAGE Lysate: 40 μg

Lane 1-2: NIH/3T3 cells human ovarian cancer tissue

Primary antibody: TA323291 (ABCE1 Antibody) at

dilution 1/300

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 1 minute