

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA323294

PSAP Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB: 1:500-2000
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
lsotype:	lgG
Clonality:	Polyclonal
Immunogen:	Fusion protein corresponding to a region derived from 60-310 amino acids of human prosaposin
Formulation:	PBS pH7.3, 0.05% NaN3, 50% glycerol
Concentration:	lot specific
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	58 kDa
Gene Name:	prosaposin
Database Link:	<u>NP_001035930</u> <u>Entrez Gene 19156 MouseEntrez Gene 25524 RatEntrez Gene 5660 Human</u> <u>P07602</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

SAP Rabbit Polyclonal Antibody – TA323294

Background:The PSAP gene encodes prosaposin, a precursor of four small nonenzymatic glycoproteins
termed 'sphingolipid activator proteins' (SAPs) that assist in the lysosomal hydrolysis of
sphingolipids. After proteolytic processing of the presaposin protein, these 4 released
polypeptides are functional activators. Saposin A is encoded by residues 60 to 143 of PSAP,
saposin B by 195 to 275, saposin C by 311 to 390, and saposin D by 405 to 487. They are four
12-14 kDa heatstable glycoproteins. Saposins A-D localize primarily to the lysosomal
compartment where they facilitate the catabolism of glycosphingolipids with short
oligosaccharide groups. Saposins A-D are required for the hydrolysis of certain sphingolipids
by specific lysosomal hydrolases. Defects in PSAP are the cause of Gaucher disease, Tay-
Sachs disease, and metachromatic leukodystrophy. This PSAP antibody (10801-1-AP) is
expected to recognize both saposin A and B.

Synonyms:	GLBA; SAP1
Protein Families:	Druggable Genome
Protein Pathways:	Lysosome

Product images:



Predicted band size: 58 kDa. Positive control: HEK-293 cell lysate. Recommended dilution: 1/500-2000. (Gel: 10%SDS-PAGE Lysate: 40 ug Primary antibody: 1/500 dilution Secondary antibody: Goat anti Rabbit IgG - H&L (HRP) at 1/10000 dilution Exposure time: 1 minute)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US