

Product datasheet for TA323304

TCF4 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications:

Recommended Dilution: WB: 1:500-2000

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: lgG

Clonality: Polyclonal

Immunogen: Fusion protein corresponding to N terminal 210 amino acids of human transcription factor 4

PBS pH7.4 0.05% NaN3, 40% glycerol Formulation:

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Store at -20°C as received. Storage:

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 71 kDa

Gene Name: transcription factor 4

Database Link: NP 001077431

Entrez Gene 21413 MouseEntrez Gene 84382 RatEntrez Gene 6925 Human

P15884

Background: This gene encodes transcription factor 4, a basic helix-loop-helix transcription factor. The

> encoded protein recognizes an Ephrussi-box ('E-box') binding site ('CANNTG') - a motif first identified in immunoglobulin enhancers. This gene is broadly expressed, and may play an important role in nervous system development. Defects in this gene are a cause of Pitt-Hopkins syndrome. In addition, an intronic CTG repeat normally numbering 10-37 repeat units can expand to >50 repeat units and cause Fuchs endothelial corneal dystrophy. Multiple alternatively spliced transcript variants that encode different proteins have been described

Synonyms: bHLHb19; E2-2; ITF-2; ITF2; PTHS; SEF-2; SEF2-1; SEF2-1A; SEF2-1B; SEF2-1D; TCF-4



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Transcription Factors

Product images:



Predicted band size: 71 kDa. Positive control: K562 cell lysate. Recommended dilution: 1/500-2000. (Gel: 8%SDS-PAGE Lysate: 40 ug Primary antibody: 1/500 dilution Secondary antibody: Goat anti Rabbit IgG - H&L (HRP) at 1/10000 dilution Exposure time: 1 minute)