

Product datasheet for TA323311S

AADAC Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Human fetal liver and liver cancer tissue

IHC: 50-100

Positive control: Human liver cancer Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein corresponding to C terminal 250 amino acids of human arylacetamide

deacetylase

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 46 kDa

Gene Name: arylacetamide deacetylase

Database Link: NP 001077

Entrez Gene 13 Human

P22760



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Microsomal arylacetamide deacetylase competes against the activity of cytosolic arylamine N-acetyltransferase; which catalyzes one of the initial biotransformation pathways for arylamine and heterocyclic amine carcinogens. Arylacetamide deacetylation is an important enzyme activity in the metabolic activation of arylamine substrates to ultimate carcinogens. Displays major serine hydrolase activity in liver microsomes. Hydrolyzes also flutamide; which is an antiandrogen drug used for the treatment of prostate cancer that occasionally causes severe hepatotoxicity. Displays cellular triglyceride lipase activity in liver. Increases intracellular fatty acids derived from hydrolysis of newly formed triglyceride stores.

Synonyms: CES5A1; DAC

Protein Families: Druggable Genome, Transmembrane

Product images:



Gel: 10%SDS-PAGE Lysate: 40 µg

Lane 1-2: Human fetal liver tissue

Human liver cancer tissue


Primary antibody: [TA323311] (AADAC Antibody)

at dilution 1/600

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 1 minute



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA323311] (AADAC Antibody) at dilution 1/30 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA323311] (AADAC Antibody) at dilution 1/30, treated with fusion protein. (Original magnification: ×200)