

Product datasheet for **TA323314**

NFkB p100 / p52 (NFKB2) Rabbit Polyclonal Antibody


Product data:

Product Type:	Primary Antibodies
Applications:	IHC
Recommended Dilution:	IHC: 50-200 Positive control: Human cervical cancer Predicted cell location: Cytoplasm
Reactivity:	Human, Mouse
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Synthetic peptide peptide corresponding to a region derived from 865-880 amino acids of human nuclear factor of kappa light polypeptide gene enhancer in B-cells 2 (p49/p100)
Formulation:	PBS pH7.3, 0.05% NaN ₃ , 50% glycerol
Concentration:	lot specific
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	nuclear factor kappa B subunit 2
Database Link:	NP_002493 Entrez Gene 18034 Mouse Entrez Gene 4791 Human Q00653




[View online »](#)

Background:	This gene encodes one of the subunits of the transcription factor complex nuclear factor-kappa-B (NFkB). The NFkB transcription factor complex is expressed in numerous cell types and functions as a central activator of genes involved in inflammation and immune function. The NFkB complex can consist of different subunits that form both homo- or heterodimers which bind specific kappa-B elements in target genes. This gene encodes the p100 subunit that is processed into the active p52 subunit. This protein can function as both a transcriptional activator and repressor; depending on its dimer partner. Alternate splicing results in both coding and non-coding variants.?
Synonyms:	CVID10; H2TF1; LYT-10; LYT10; NF-kB2; p52; p100
Protein Families:	Transcription Factors
Protein Pathways:	MAPK signaling pathway, Pathways in cancer

Product images:

Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA323314 (NFKB2 Antibody) at dilution 1/20 (Original magnification: x200)



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA323314 (NFKB2 Antibody) at dilution 1/20, treated with synthetic peptide. (Original magnification: x200)