

## **Product datasheet for TA323326S**

## **NLRP3 Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 15-50

Positive control: Human esophagus cancer

Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Fusion protein corresponding to a region derived from 1-301 amino acids of human NLR

family, pyrin domain containing 3

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Gene Name: NLR family, pyrin domain containing 3

Database Link: NP 899632

Entrez Gene 114548 Human

Q96P20



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:


This gene encodes a pyrin-like protein containing a pyrin domain, a nucleotide-binding site (NBS) domain, and a leucine-rich repeat (LRR) motif. This protein interacts with the apoptosis-associated speck-like protein PYCARD/ASC, which contains a caspase recruitment domain, and is a member of the NALP3 inflammasome complex. This complex functions as an upstream activator of NF-kappaB signaling, and it plays a role in the regulation of inflammation, the immune response, and apoptosis. Mutations in this gene are associated with familial cold autoinflammatory syndrome (FCAS), Muckle-Wells syndrome (MWS), chronic infantile neurological cutaneous and articular (CINCA) syndrome, and neonatal-onset multisystem inflammatory disease (NOMID). Multiple alternatively spliced transcript variants encoding distinct isoforms have been identified for this gene. Alternative 5' UTR structures are suggested by available data; however, insufficient evidence is available to determine if all of the represented 5' UTR splice patterns are biologically valid.?

Synonyms: AGTAVPRL; AII; AVP; C1orf7; CIAS1; CLR1.1; FCAS; FCAS1; FCU; MWS; NALP3; PYPAF1

**Protein Families:** Druggable Genome


**Protein Pathways:** NOD-like receptor signaling pathway

## **Product images:**




Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using [TA323326] (NLRP3 Antibody) at dilution 1/15 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using [TA323326] (NLRP3 Antibody) at dilution 1/15, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA323326] (NLRP3 Antibody) at dilution 1/15 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA323326] (NLRP3 Antibody) at dilution 1/15, treated with fusion protein. (Original magnification: ×200)