

## **Product datasheet for TA323344**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436

OriGene Technologies, Inc.

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## 5HT4 Receptor (HTR4) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB: 200-1000

WB positive control: Hela, 293T, A549 and Jurkat cells

**Reactivity:** Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Synthetic peptide corresponding to a region derived from 45-59 amino acids of Human 5-

hydroxytryptamine (serotonin) receptor 4, G protein-coupled

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 44 kDa

**Gene Name:** 5-hydroxytryptamine receptor 4

Database Link: NP 001035263

Entrez Gene 15562 MouseEntrez Gene 25324 RatEntrez Gene 3360 Human

Q13639

**Background:** This gene is a member of the family of serotonin receptors; which are G protein coupled

receptors that stimulate cAMP production in response to serotonin (5-hydroxytryptamine). The gene product is a glycosylated transmembrane protein that functions in both the

peripheral and central nervous system to modulate the release of various neurotransmitters. Multiple transcript variants encoding proteins with distinct C-terminal sequences have been

described.

**Synonyms:** 5-HT4; 5-HT4R






**Protein Families:** Druggable Genome, GPCR, Transmembrane

**Protein Pathways:** Calcium signaling pathway, Neuroactive ligand-receptor interaction

## **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 µg Lane 1-4: Hela cells 293T cells

293T cells A549 cells Jurkat cells

Primary antibody: TA323344 (HTR4 Antibody) at

dilution 1/200

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 3 minutes