

Product datasheet for **TA323529S**

ACTH (POMC) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB: 500-2000 WB positive control: Human fetal brain tissue
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Synthetic peptide peptide corresponding to a region derived from 136-150 amino acids of human proopiomelanocortin
Formulation:	PBS pH7.3, 0.05% NaN ₃ , 50% glycerol
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	29 kDa
Gene Name:	proopiomelanocortin
Database Link:	NP_000930 Entrez Gene 18976 Mouse Entrez Gene 24664 Rat Entrez Gene 5443 Human P01189



[View online »](#)

Background:

This gene encodes a polypeptide hormone precursor that undergoes extensive; tissue-specific; post-translational processing via cleavage by subtilisin-like enzymes known as prohormone convertases. There are eight potential cleavage sites within the polypeptide precursor and; depending on tissue type and the available convertases; processing may yield as many as ten biologically active peptides involved in diverse cellular functions. The encoded protein is synthesized mainly in corticotroph cells of the anterior pituitary where four cleavage sites are used; adrenocorticotrophin; essential for normal steroidogenesis and the maintenance of normal adrenal weight; and lipotropin beta are the major end products. In other tissues; including the hypothalamus; placenta; and epithelium; all cleavage sites may be used; giving rise to peptides with roles in pain and energy homeostasis; melanocyte stimulation; and immune modulation.

Synonyms:


ACTH; CLIP; LPH; MSH; NPP; POC

Protein Families:

Druggable Genome

Protein Pathways:

Adipocytokine signaling pathway, Melanogenesis

Product images:

Gel: 10%SDS-PAGE

Lysate: 40 µg

Lane: Human fetal brain tissue

Primary antibody: [TA323529] (POMC(Alpha-MSH) Antibody) at dilution 1/400

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 10 seconds