

Product datasheet for TA323531

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Nociceptin receptor (OPRL1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 500-2000

WB positive control: Human liver cancer tissue

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide corresponding to a region derived from 360-380 amino acids of Human

opiate receptor-like 1

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 41 kDa

Gene Name: opioid related nociceptin receptor 1

Database Link: NP 000904

Entrez Gene 18389 MouseEntrez Gene 29256 RatEntrez Gene 4987 Human

P41146

Background: The protein encoded by this gene is a G protein-coupled receptor whose expression can be

induced by phytohemagglutinin. The encoded integral membrane protein is a receptor for the 17 aa neuropeptide nociceptin/orphanin FQ. This gene may be involved in the regulation of numerous brain activities; particularly instinctive and emotional behaviors. A promoter for this gene also functions as a promoter for another gene; regulator of G-protein signalling 19 (RGS19); located on the opposite strand. Three transcript variants encoding the same protein

have been found for this gene.




Nociceptin receptor (OPRL1) Rabbit Polyclonal Antibody - TA323531

Synonyms: KOR-3; NOCIR; NOP; NOPr; OOR; ORL1

Protein Families: Druggable Genome, GPCR, Transmembrane

Protein Pathways: Neuroactive ligand-receptor interaction

Product images:



Gel: 10%SDS-PAGE Lysate: 40 μg

Lane: Human liver cancer tissue

Primary antibody: TA323531 (OPRL1 Antibody) at

dilution 1/900

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 1 minute