

Product datasheet for **TA323551**

NMDAR2B (GRIN2B) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB: 500-2000 WB positive control: Mouse brain tissue
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Synthetic peptide corresponding to a region derived from 1413-1425 amino acids of Human glutamate receptor, ionotropic, N-methyl D-aspartate 2B
Formulation:	PBS pH7.3, 0.05% NaN ₃ , 50% glycerol
Concentration:	lot specific
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	166 kDa
Gene Name:	glutamate ionotropic receptor NMDA type subunit 2B
Database Link:	NP_000825 Entrez Gene 14812 Mouse Entrez Gene 24410 Rat Entrez Gene 2904 Human Q13224
Background:	N-methyl-D-aspartate (NMDA) receptors are a class of ionotropic glutamate receptors. NMDA receptor channel has been shown to be involved in long-term potentiation; an activity-dependent increase in the efficiency of synaptic transmission thought to underlie certain kinds of memory and learning. NMDA receptor channels are heteromers composed of three different subunits: NR1 (GRIN1); NR2 (GRIN2A; GRIN2B; GRIN2C; or GRIN2D) and NR3 (GRIN3A or GRIN3B). The NR2 subunit acts as the agonist binding site for glutamate. This receptor is the predominant excitatory neurotransmitter receptor in the mammalian brain.?



[View online »](#)

Synonyms: EIEE27; GluN2B; hNR3; MRD6; NMDAR2B; NR2B

Protein Families: Druggable Genome, Ion Channels: Glutamate Receptors, Transmembrane

Protein Pathways: Alzheimer's disease, Amyotrophic lateral sclerosis (ALS), Huntington's disease, Long-term potentiation, Neuroactive ligand-receptor interaction, Systemic lupus erythematosus

Product images:



Gel: 8%SDS-PAGE
Lysate: 40 µg
Lane: Mouse brain tissue
Primary antibody: TA323551 (GRIN2B Antibody)
at dilution 1/700
Secondary antibody: Goat anti rabbit IgG at
1/8000 dilution
Exposure time: 40 seconds