

## **Product datasheet for TA323558S**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Ionotropic Glutamate receptor 2 (GRIA2) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 100-300

Positive control: Human liver cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Synthetic peptide corresponding to a region derived from 263-276 amino acids of Human

glutamate receptor, ionotropic, AMPA 2

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** glutamate ionotropic receptor AMPA type subunit 2

Database Link: NP 000817

Entrez Gene 14800 MouseEntrez Gene 29627 RatEntrez Gene 2891 Human

P42262





Background:

Glutamate receptors are the predominant excitatory neurotransmitter receptors in the mammalian brain and are activated in a variety of normal neurophysiologic processes. This gene product belongs to a family of glutamate receptors that are sensitive to alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionate (AMPA); and function as ligand-activated cation channels. These channels are assembled from 4 related subunits; GRIA1-4. The subunit encoded by this gene (GRIA2) is subject to RNA editing (CAG->CGG; Q->R) within the second transmembrane domain; which is thought to render the channel impermeable to Ca(2+). Human and animal studies suggest that pre-mRNA editing is essential for brain function; and defective GRIA2 RNA editing at the Q/R site may be relevant to amyotrophic lateral sclerosis (ALS) etiology. Alternative splicing; resulting in transcript variants encoding different isoforms; (including the flip and flop isoforms that vary in their signal transduction properties); has been noted for this gene.


**Synonyms:** GluA2; GluR-K2; GLUR2; GLURB; HBGR2

**Protein Families:** Druggable Genome, Ion Channels: Glutamate Receptors, Transmembrane

Protein Pathways: Amyotrophic lateral sclerosis (ALS), Long-term depression, Long-term potentiation,


Neuroactive ligand-receptor interaction

## **Product images:**




Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA323558] (GRIA2 Antibody) at dilution 1/70 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA323558] (GRIA2 Antibody) at dilution 1/70, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human colon cancer tissue using [TA323558] (GRIA2 Antibody) at dilution 1/70 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human colon cancer tissue using [TA323558] (GRIA2 Antibody) at dilution 1/70, treated with synthetic peptide. (Original magnification: ×200)