

Product datasheet for TA323595

Aminoacylase 1 (ACY1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies Applications: IHC, WB Recommended Dilution: WB: 200-1000 WB positive control: Mouse kidney and human normal kidney tissue, K562 cells IHC: 25-100 Positive control: Human thyroid cancer Predicted cell location: Cytoplasm and Nucleus **Reactivity:** Human, Mouse Host: Rabbit Isotype: lgG **Clonality:** Polyclonal Immunogen: Fusion protein corresponding to C terminal 250 amino acids of human aminoacylase 1 PBS pH7.3, 0.05% NaN3, 50% glycerol Formulation: Concentration: lot specific **Purification:** Antigen affinity purification **Conjugation:** Unconjugated Storage: Store at -20°C as received. Stable for 12 months from date of receipt. Stability: **Predicted Protein Size:** 46 kDa Gene Name: aminoacylase 1 Database Link: NP 000657 Entrez Gene 109652 MouseEntrez Gene 95 Human Q03154



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.


9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GRIGENE Aminoacylase 1 (ACY1) Rabbit Polyclonal Antibody – TA323595

Background:This gene encodes a cytosolic, homodimeric, zinc-binding enzyme that catalyzes the
hydrolysis of acylated L-amino acids to L-amino acids and an acyl group, and has been
postulated to function in the catabolism and salvage of acylated amino acids. This gene is
located on chromosome 3p21.1, a region reduced to homozygosity in small-cell lung cancer
(SCLC), and its expression has been reported to be reduced or undetectable in SCLC cell lines
and tumors. The amino acid sequence of human aminoacylase-1 is highly homologous to the
porcine counterpart, and this enzyme is the first member of a new family of zinc-binding
enzymes. Mutations in this gene cause aminoacylase-1 deficiency, a metabolic disorder
characterized by central nervous system defects and increased urinary excretion of N-
acetylated amino acids. Alternative splicing of this gene results in multiple transcript variants.
Read-through transcription also exists between this gene and the upstream ABHD14A
(abhydrolase domain containing 14A) gene, as represented in GeneID:100526760. A related
pseudogene has been identified on chromosome 18.


Synonyms:	ACY-1; ACY1D; HEL-S-5
Protein Families:	Protease
Protein Pathways:	Arginine and proline metabolism, Metabolic pathways

Product images:



Gel: 10%SDS-PAGE Lysate: 40 µg Lane 1-3: Mouse kidney tissue human normal kidney tissue K562 cells Primary antibody: TA323595 (ACY1 Antibody) at dilution 1/200 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution Exposure time: 1 minute

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA323595 (ACY1 Antibody) at dilution 1/20 (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA323595 (ACY1 Antibody) at dilution 1/20, treated with fusion protein. (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US