

## **Product datasheet for TA323596**

### **OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

# Alcohol Dehydrogenase (ADH1A) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 25-100

Positive control: Human gasrtic cancer Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Fusion protein corresponding to C terminal 300 amino acids of human alcohol

dehydrogenase 1A (class I), alpha polypeptide

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** alcohol dehydrogenase 1A (class I), alpha polypeptide

Database Link: NP 000658

Entrez Gene 124 Human

P07327

#### Alcohol Dehydrogenase (ADH1A) Rabbit Polyclonal Antibody - TA323596

**Background:** This gene encodes a member of the alcohol dehydrogenase family. The encoded protein is

the alpha subunit of class I alcohol dehydrogenase, which consists of several homo- and heterodimers of alpha, beta and gamma subunits. Alcohol dehydrogenases catalyze the oxidation of alcohols to aldehydes. This gene is active in the liver in early fetal life but only weakly active in adult liver. This gene is found in a cluster with six additional alcohol dehydrogenase genes, including those encoding the beta and gamma subunits, on the long arm of chromosome 4. Mutations in this gene may contribute to variation in certain

personality traits and substance dependence.

Synonyms: ADH1


**Protein Families:** Druggable Genome

**Protein Pathways:** Drug metabolism - cytochrome P450, Fatty acid metabolism, Glycolysis / Gluconeogenesis,

Metabolic pathways, Metabolism of xenobiotics by cytochrome P450, Retinol metabolism,


Tyrosine metabolism

# **Product images:**




Immunohistochemistry of paraffin-embedded Human gasrtic cancer tissue using TA323596 (ADH1A Antibody) at dilution 1/25 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human gasrtic cancer tissue using TA323596 (ADH1A Antibody) at dilution 1/25, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA323596 (ADH1A Antibody) at dilution 1/25 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA323596 (ADH1A Antibody) at dilution 1/25, treated with fusion protein. (Original magnification: ×200)