

Product datasheet for TA323688S

TAT Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: Human normal liver tissue

IHC: 25-100

Positive control: Human colon cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein corresponding to a region derived from C terminal 250 amino acids of human

tyrosine aminotransferase

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: tyrosine aminotransferase

Database Link: NP 000344

Entrez Gene 24813 RatEntrez Gene 234724 MouseEntrez Gene 6898 Human

P17735

Background: This nuclear gene encodes a mitochondrial protein tyrosine aminotransferase which is

present in the liver and catalyzes the conversion of L-tyrosine into p-hydroxyphenylpyruvate. Mutations in this gene cause tyrosinemia (type II; Richner-Hanhart syndrome); a disorder accompanied by major skin and corneal lesions; with possible mental retardation. A regulator

gene for tyrosine aminotransferase is X-linked.

Synonyms: cytosolic; tyrosine aminotransferase



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Protein Families: Druggable Genome, Embryonic stem cells, ES Cell Differentiation/IPS

Protein Pathways: Cysteine and methionine metabolism, Metabolic pathways, Phenylalanine, tyrosine and

 $tryptophan\ biosynthesis,\ Phenylalanine\ metabolism,\ Tyrosine\ metabolism,\ Ubiquinone\ and$

other terpenoid-quinone biosynthesis

Product images:



Gel: 8%SDS-PAGE Lysate: 40 μg

Lane: Human normal liver tissue


Primary antibody: [TA323688] (TAT Antibody) at

dilution 1/200

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 3 minutes




Immunohistochemistry of paraffin-embedded Human colon cancer tissue using [TA323688] (TAT Antibody) at dilution 1/35 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human colon cancer tissue using [TA323688] (TAT Antibody) at dilution 1/35, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA323688] (TAT Antibody) at dilution 1/35 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA323688] (TAT Antibody) at dilution 1/35, treated with fusion protein. (Original magnification: ×200)