

## **Product datasheet for TA323812**

## p73 (TP73) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: HepG2, Hela, K562 cell lysates

IHC: 50-200

Positive control: Human esophagus cancer

Predicted cell location: Nucleus

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

**Clonality:** Polyclonal

**Immunogen:** Fusion protein corresponding to a region derived from 131-310 amino acids of human tumor

protein p73

**Formulation:** PBS pH7.3, 0.05% NaN3, 50% glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 70 kDa

**Gene Name:** tumor protein p73

Database Link: NP 001119712

Entrez Gene 7161 Human

O15350



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:


This gene encodes a member of the p53 family of transcription factors involved in cellular responses to stress and development. It maps to a region on chromosome 1p36 that is frequently deleted in neuroblastoma and other tumors; and thought to contain multiple tumor suppressor genes. The demonstration that this gene is monoallelically expressed (likely from the maternal allele); supports the notion that it is a candidate gene for neuroblastoma. Many transcript variants resulting from alternative splicing and/or use of alternate promoters have been found for this gene; but the biological validity and the full-length nature of some variants have not been determined.

Synonyms: P73

**Protein Families:** Druggable Genome, Transcription Factors

**Protein Pathways:** Neurotrophin signaling pathway, p53 signaling pathway

## **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 µg Lane 1-3: HepG2 Hela

K562 cell lysates


Primary antibody: TA323812 (TP73 Antibody) at

dilution 1/550

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 1 minute




Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA323812 (TP73 Antibody) at dilution 1/70 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA323812 (TP73 Antibody) at dilution 1/70, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA323812 (TP73 Antibody) at dilution 1/70 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA323812 (TP73 Antibody) at dilution 1/70, treated with fusion protein. (Original magnification: ×200)