

Product datasheet for TA323841

Froduct datasireet for TA32304

EIF4E Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IF, IHC, WB

Recommended Dilution: WB: 1:500-1000, IHC: 1:50-100, IF: 1:100-200

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Peptide sequence around aa. 207~211 (S-G-S-T-T) derived from Human eIF4E.

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 25 kDa

Gene Name: eukaryotic translation initiation factor 4E

Database Link: NP 001124150

Entrez Gene 13684 MouseEntrez Gene 117045 RatEntrez Gene 1977 Human

P06730

Background: Recognizes and binds the 7-methylguanosine-containing mRNA cap during an early step in

the initiation of protein synthesis and facilitates ribosome binding by inducing the unwinding

of the mRNAs secondary structures.

Synonyms: AUTS19; CBP; eIF-4E; EIF4E1; EIF4EL1; EIF4F

Protein Pathways: Insulin signaling pathway, mTOR signaling pathway




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Product images:



Predicted band size: 25 kDa. Positive control: Hela, HepG2 and 293 cells lysate. Recommended dilution: 1/500-1000. (Gel: 10%SDS-PAGE Lane 1, 2, 3: Hela, HepG2 and 293 cells Lysates: 30 ug per lane Primary antibody: 1/500 dilution Secondary antibody: Goat anti Rabbit IgG - H&L (HRP) at 1/10000 dilution Exposure time: 1 minute)



Predicted cell location: Cytoplasm. Positive control: Human breast carcinoma tissue. Recommended dilution: 1/50-100 The image on the left is immunohistochemistry of paraffinembedded human breast carcinoma tissue using EIF4E (Ab-209) antibody at dilution 1/50, on the right is treated with the synthetic peptide. (Original magnification: x200)



Predicted cell location: Cytoplasm. Positive control: Hela cells. Recommended dilution: 1/100-200. The image is immunofluorescence of methanol-fixed Hela cells using EIF4E antibody at dilution 1/100. (Original magnification: ×200)