

Product datasheet for TA323894

OGA Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Mouse skin tissue lysate

IHC: 100-300

Positive control: Human colorectal cancer Predicted cell location: Cytoplasm and Nucleus

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein corresponding to a region derived from 695-814 amino acids of human

meningioma expressed antigen 5 (hyaluronidase)

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 103 kDa

Gene Name: meningioma expressed antigen 5 (hyaluronidase)

Database Link: NP 001135906

Entrez Gene 76055 MouseEntrez Gene 154968 RatEntrez Gene 10724 Human

<u>O60502</u>



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

The dynamic modification of cytoplasmic and nuclear proteins by O-linked N-acetylglucosamine (O-GlcNAc) addition and removal on serine and threonine residues is catalyzed by OGT; which adds O-GlcNAc; and MGEA5; a glycosidase that removes O-GlcNAc modifications. Cleaves GlcNAc but not GalNAc from glycopeptides. Can use p-nitrophenyl-beta-GlcNAc as substrate but not p-nitrophenyl-beta-GalNAc or p-nitrophenyl-alpha-GlcNAc. Possesses hyaluronidase activity. Acetylates 'Lys-8' of histone H4 and 'Lys-14' of histone H3. Shows highest expression in the brain; placenta and pancreas.

Synonyms: MEA5; NCOAT; OGA

Product images:



Gel: 6%SDS-PAGE Lysate: 40 μg

Lane: Mouse skin tissue lysate


Primary antibody: TA323894 (OGA Antibody) at

dilution 1/400

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 1 minute




Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using TA323894 (OGA Antibody) at dilution 1/60 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using TA323894 (OGA Antibody) at dilution 1/60, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA323894 (OGA Antibody) at dilution 1/60 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA323894 (OGA Antibody) at dilution 1/60, treated with fusion protein. (Original magnification: ×200)