

Product datasheet for TA323908S

BDNF Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: Mouse skin tissue

IHC: 10-50

Positive control: Human colon cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein corresponding to a region derived from 19-247 amino acids of human brain-

derived neurotrophic factor

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 28 kDa

Gene Name: brain-derived neurotrophic factor

Database Link: NP 001137277

Entrez Gene 12064 MouseEntrez Gene 24225 RatEntrez Gene 627 Human

P23560

Background: The protein encoded by this gene is a member of the nerve growth factor family. It is induced

by cortical neurons; and is necessary for survival of striatal neurons in the brain. Expression of this gene is reduced in both Alzheimer's and Huntington disease patients. This gene may play a role in the regulation of stress response and in the biology of mood disorders. Multiple

transcript variants encoding distinct isoforms have been described for this gene.?



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms: ANON2; BULN2

Protein Families: Adult stem cells, Druggable Genome, Embryonic stem cells, ES Cell Differentiation/IPS,

Induced pluripotent stem cells, Secreted Protein, Transmembrane

Protein Pathways: Huntington's disease, MAPK signaling pathway, Neurotrophin signaling pathway

Product images:



Gel: 10%SDS-PAGE Lysate: 40 µg

Lane: Mouse skin tissue


Primary antibody: [TA323908] (BDNF Antibody) at

dilution 1/250

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 2 minutes



Immunohistochemistry of paraffin-embedded Human colon cancer tissue using [TA323908] (BDNF Antibody) at dilution 1/20 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human colon cancer tissue using [TA323908] (BDNF Antibody) at dilution 1/20, treated with fusion protein. (Original magnification: ×200)