

Product datasheet for TA323924

uPA (PLAU) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 50-200

Positive control: Human liver cancer Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein corresponding to a region derived from 179-417 amino acids of human

plasminogen activator, urokinase

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: plasminogen activator, urokinase

Database Link: NP 002649

Entrez Gene 5328 Human

P00749



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:


This gene encodes a serine protease involved in degradation of the extracellular matrix and possibly tumor cell migration and proliferation. A specific polymorphism in this gene may be associated with late-onset Alzheimer's disease and also with decreased affinity for fibrin-binding. This protein converts plasminogen to plasmin by specific cleavage of an Arg-Val bond in plasminogen. Plasmin in turn cleaves this protein at a Lys-lle bond to form a two-chain derivative in which a single disulfide bond connects the amino-terminal A-chain to the catalytically active; carboxy-terminal B-chain. This two-chain derivative is also called HMW-uPA (high molecular weight uPA). HMW-uPA can be further processed into LMW-uPA (low molecular weight uPA) by cleavage of chain A into a short chain A (A1) and an amino-terminal fragment. LMW-uPA is proteolytically active but does not bind to the uPA receptor. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Synonyms: ATF; BDPLT5; QPD; u-PA; UPA; URK

Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Protease


Protein Pathways: Complement and coagulation cascades

Product images:




Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA323924 (PLAU Antibody) at dilution 1/90 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA323924 (PLAU Antibody) at dilution 1/90, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA323924 (PLAU Antibody) at dilution 1/90 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA323924 (PLAU Antibody) at dilution 1/90, treated with fusion protein. (Original magnification: ×200)